arXiv:1207.0071v1 [nlin.SI] 30 Jun 2012

The anisotropic Higgs oscillator on the two-dimensional sphere and the hyperbolic plane

Ángel Ballesteros[1], Francisco J. Herranz[1] and Fabio Musso[2]

[1] Departamento de Física, Universidad de Burgos, E-09001 Burgos, Spain

[2] Dipartimento di Fisica "Edoardo Amaldi", Universitá Roma Tre, I-00146 Rome, Italy

E-mail: `angelb@ubu.es`, `fjherranz@ubu.es`, `fmusso@ubu.es`

Abstract

An integrable generalization on the two-dimensional sphere $\mathbf{S}^2$ and the hyperbolic plane $\mathbf{H}^2$ of the Euclidean anisotropic oscillator Hamiltonian with "centrifugal" terms given by

$$\mathcal{H} = \frac{1}{2}(p_1^2 + p_2^2) + \delta q_1^2 + (\delta + \Omega) q_2^2 + \frac{\lambda_1}{q_1^2} + \frac{\lambda_2}{q_2^2}$$

is presented. The resulting generalized Hamiltonian $\mathcal{H}_\kappa$ depends explicitly on the Gaussian curvature κ of the underlying space, in such a way that all the results here presented hold simultaneously for $\mathbf{S}^2$ ($\kappa > 0$), $\mathbf{H}^2$ ($\kappa < 0$) and $\mathbf{E}^2$ ($\kappa = 0$). Moreover, $\mathcal{H}_\kappa$ is explicitly shown to be integrable (albeit not superintegrable) for any values of the parameters δ, Ω, λ_1 and λ_2. Therefore, $\mathcal{H}_\kappa$ can also be interpreted as an anisotropic generalization of the curved Higgs oscillator, that is recovered as the isotropic limit $\Omega \to 0$ of $\mathcal{H}_\kappa$. Furthermore, numerical integration of some of the trajectories for $\mathcal{H}_\kappa$ are worked out and the dynamical features arising from the introduction of a curved background are highlighted. In particular we focus on the case $\Omega = 3\delta$, whose Euclidean limit $\kappa \to 0$ is the superintegrable 1:2 oscillator. In this case we illustrate how the Euclidean superintegrability of $\mathcal{H}$ is broken when the curvature κ is non-zero. Since for the specific $\Omega = 3\delta$ case another superintegrable curved generalization is already known in the literature, the existence of an Ω-dependent plurality of integrable curved systems whose Euclidean limit is the same anisotropic oscillator $\mathcal{H}$ can be conjectured. Finally, the geometric interpretation of the curved "centrifugal" terms appearing in $\mathcal{H}_\kappa$ is also discussed in detail.

PACS: 02.30.Ik 02.40.Ky

KEYWORDS: anisotropic oscillator, integrable systems, Lie algebras, Lie–Poisson algebras, Gaussian curvature, sphere, hyperbolic space, integrable deformation, Higgs oscillator

1 Introduction

The two-dimensional (2D) anisotropic oscillator with “centrifugal” (or “Rosochatius”) terms

$$\mathcal{H} = \frac{1}{2}(p_1^2 + p_2^2) + \delta q_1^2 + (\delta + \Omega) q_2^2 + \frac{\lambda_1}{q_1^2} + \frac{\lambda_2}{q_2^2}, \tag{1.1}$$

with δ, Ω and λ_i being real parameters and $\{q_i, p_j\} = \delta_{ij}$ $(i, j = 1, 2)$, is one of the most elementary (albeit instructive) examples of completely integrable Hamiltonian systems. Its complete integrability (in the Liouville sense) is a straightforward consequence of its manifest separability in Cartesian coordinates, and for some specific choices of the parameters the system turns out to be superintegrable due to the existence of an additional independent integral of the motion.

The aim of this paper is to present an integrable generalization of this Hamiltonian on the 2D sphere $\mathbf{S}^2$ and on the hyperbolic plane $\mathbf{H}^2$, in the sense that the Hamiltonian (1.1) will be recovered when the Euclidean limit is considered. In fact, throughout the paper we shall make use of the constant Gaussian curvature κ of these two curved spaces as an explicit parameter, and all the results here presented will be simultaneously valid for $\mathbf{S}^2$ $(\kappa > 0)$, $\mathbf{H}^2$ $(\kappa < 0)$ and $\mathbf{E}^2$ $(\kappa = 0)$. In this way, the “flat/linear” limit $\kappa \to 0$ in the spherical and hyperbolic systems will be always well defined leading to the corresponding Euclidean Hamiltonian (1.1) (obviously, the term linear refers to the equations of motion when both λ_i vanish). Conversely, the non-trivial dynamical nature of the transition from the “flat/linear” systems defined on $\mathbf{E}^2$ to the “curved/nonlinear” ones associated to $\mathbf{S}^2$ and $\mathbf{H}^2$ can be clearly appreciated through this approach.

Since integrability will be our essential guiding principle when constructing the curved generalization of the system, let us briefly recall the (Liouville) integrability properties of (1.1) in terms of the δ, Ω and λ_i parameters. Obviously, $\mathcal{H}$ is always integrable since it is separable in Cartesian coordinates:

$$\mathcal{H} = \mathcal{I}_1 + \mathcal{I}_2, \qquad \mathcal{I}_1 = \frac{1}{2}p_1^2 + \delta q_1^2 + \frac{\lambda_1}{q_1^2}, \qquad \mathcal{I}_2 = \frac{1}{2}p_2^2 + (\delta + \Omega) q_2^2 + \frac{\lambda_2}{q_2^2}. \tag{1.2}$$

Thus each pair $(\mathcal{H}, \mathcal{I}_1)$ and $(\mathcal{H}, \mathcal{I}_2)$ is formed by two functionally independent functions in involution with respect to the canonical Poisson bracket. However, there are two particular situations for which (1.1) is superintegrable:

- The isotropic oscillator $(\Omega = 0)$ with Rosochatius terms (λ_i arbitrary), which is frequently called the Smorodinsky–Winternitz (SW) system [1, 2, 3, 4]. In this case the additional integral of the motion is quadratic in the momenta.
- The anisotropic oscillator $(\Omega \neq 0)$ with Rosochatius terms (λ_i arbitrary) provided that the ratio of the associated frequencies is a rational number [5, 6]. In this family of systems only the commensurate 1:2 oscillator $(\Omega = 3\delta)$ plus a *single* Rosochatius potential (say $\lambda_2 = 0$) presents an additional integral which is quadratic in the momenta (if both $\lambda_i \neq 0$ or the anisotropic oscillator has any other pair of commensurate frequencies the additional integral is of higher order in the momenta [5, 6]). The resulting 1:2

superintegrable Hamiltonian and its quadratic integrals of motion turn out to be

$$\begin{aligned}
&\mathcal{H}' = \frac{1}{2}(p_1^2 + p_2^2) + \delta q_1^2 + 4\delta q_2^2 + \frac{\lambda_1}{q_1^2}, \qquad \mathcal{H}' = \mathcal{I}_1' + \mathcal{I}_2', \\
&\mathcal{I}_1' \equiv \mathcal{I}_1 = \frac{1}{2}p_1^2 + \delta q_1^2 + \frac{\lambda_1}{q_1^2}, \qquad \mathcal{I}_2' = \frac{1}{2}p_2^2 + 4\delta q_2^2, \\
&\mathcal{L}' = (q_1 p_2 - q_2 p_1)p_1 + 2\delta q_1^2 q_2 - \frac{2\lambda_1 q_2}{q_1^2}.
\end{aligned} \tag{1.3}$$

It can be straightforwardly checked that the sets $(\mathcal{H}', \mathcal{I}_1', \mathcal{L}')$ and $(\mathcal{H}', \mathcal{I}_2', \mathcal{L}')$ are formed by three functionally independent functions.

To the best of our knowledge, there are only two particular cases of the Euclidean oscillator $\mathcal{H}$ (1.1) whose integrable generalization to the 2D sphere and hyperbolic plane are known. The first one is a curved generalization of the isotropic $\Omega = 0$ oscillator, which is the so-called Higgs oscillator [7, 8]; in this case when both centrifugal terms are taken different from zero the resulting Hamiltonian provides the curved SW system [9, 10, 11, 12, 13]. The second one is a curved analogue $\mathcal{H}_\kappa'$ of the superintegrable nonisotropic 1:2 system $\mathcal{H}'$ (1.3), which was found in [11]. Remarkably enough, these two curved systems are also superintegrable.

Therefore, a "generic" (for any value of Ω) integrable generalization of $\mathcal{H}$ (1.1) on $\mathbf{S}^2$ and $\mathbf{H}^2$ was still lacking, and we present it in this paper. We will denote it $\mathcal{H}_\kappa$ and, since the limit $\Omega \to 0$ of $\mathcal{H}_\kappa$ will lead to the Higgs oscillator, we shall call this new system the "anisotropic Higgs oscillator".

We would like to emphasize that $\mathcal{H}_\kappa$ seems to be superintegrable *only in the* $\Omega = 0$ *case* (i.e., the isotropic Higgs oscillator or, equivalently, the 1:1 case). In fact, numerical analysis of bounded trajectories for $\mathcal{H}_\kappa$ with $\Omega = 3\delta$ shows that they are not periodic, in contrast with what it should be expected for any superintegrable system. Moreover we find that the curved analogue of $\mathcal{H}'$, i.e. $\mathcal{H}_\kappa'$, is no longer a particular case of $\mathcal{H}_\kappa$ with $\Omega = 3\delta$ and $\lambda_2 = 0$, which proves that both Hamiltonians define two *different* curved systems. On the contrary, bounded trajectories of $\mathcal{H}_\kappa'$ are found to be periodic in agreement with its superintegrability nature. All these results suggest that the curved superintegrable analogues of different commensurate Euclidean oscillators could be given by different systems that would "collapse" to a common form in the Euclidean limit $\kappa \to 0$.

The paper is organized as follows. In the next section we briefly recall the basics on Beltrami and geodesic polar variables that generalize the Cartesian and polar ones, respectively, to $\mathbf{S}^2$ and $\mathbf{H}^2$. In section 3 we introduce the anisotropic Higgs oscillator $\mathcal{H}_\kappa$ and we give the explicit form of its integral of the motion for generic values of the parameters. We also present some numerical analysis of its trajectories for some commensurate values of its Euclidean frequencies, showing that superintegrability does not survive under the deformation introduced through the curvature. In section 4 we analyse the superintegrable curved oscillator given by $\mathcal{H}_\kappa'$, and the 1:2 Lissajous curves on $\mathbf{S}^2$ and $\mathbf{H}^2$ are found as the trajectories for the system. In section 5 we perform a detailed geometrical analysis of the "curved" centrifugal terms [14] of all the previous curved oscillators, and their dynamical effects are numerically explored. Finally, the last section is devoted to some concluding remarks and open problems.

2 Geodesic polar and Beltrami variables

First of all, let us rewrite the Euclidean Hamiltonians given in the previous section in terms of polar coordinates (r, ϕ) with conjugate momenta (p_r, p_ϕ). This change of coordinates induces the following canonical transformation

$$q_1 = r\cos\phi, \quad q_2 = r\sin\phi, \quad p_1 = \cos\phi\, p_r - \frac{\sin\phi}{r}\, p_\phi, \quad p_2 = \sin\phi\, p_r + \frac{\cos\phi}{r}\, p_\phi, \tag{2.1}$$

that transforms the expressions (1.1)–(1.3) into

$$\begin{aligned}
\mathcal{H} &= \frac{1}{2}\left(p_r^2 + \frac{p_\phi^2}{r^2}\right) + \delta r^2 + \Omega\, r^2 \sin^2\phi + \frac{\lambda_1}{r^2\cos^2\phi} + \frac{\lambda_2}{r^2\sin^2\phi}, \\
\mathcal{I}_1 &= \frac{1}{2}\left(\cos\phi\, p_r - \frac{\sin\phi}{r}\, p_\phi\right)^2 + \delta r^2\cos^2\phi + \frac{\lambda_1}{r^2\cos^2\phi}, \\
\mathcal{I}_2 &= \frac{1}{2}\left(\sin\phi\, p_r + \frac{\cos\phi}{r}\, p_\phi\right)^2 + (\delta+\Omega) r^2\sin^2\phi + \frac{\lambda_2}{r^2\sin^2\phi};
\end{aligned} \tag{2.2}$$

$$\begin{aligned}
\mathcal{H}' &= \frac{1}{2}\left(p_r^2 + \frac{p_\phi^2}{r^2}\right) + \delta r^2 + 3\delta\, r^2\sin^2\phi + \frac{\lambda_1}{r^2\cos^2\phi}, \\
\mathcal{I}'_1 &\equiv \mathcal{I}_1, \qquad \mathcal{I}'_2 = \frac{1}{2}\left(\sin\phi\, p_r + \frac{\cos\phi}{r}\, p_\phi\right)^2 + 4\delta r^2\sin^2\phi, \\
\mathcal{L}' &= \left(\cos\phi\, p_r - \frac{\sin\phi}{r}\, p_\phi\right) p_\phi + 2\delta r^3\cos^2\phi\sin\phi - \frac{2\lambda_1\tan\phi}{r\cos\phi}.
\end{aligned} \tag{2.3}$$

Now, in order to generalize the two above *flat* systems (2.2) and (2.3) to the sphere and to the hyperbolic plane we will apply a geometric aproach in which $\mathbf{S}^2$ and $\mathbf{H}^2$ are obtained as homogeneous spaces of certain 3D Lie groups. In this way, the realization of the associated symmetry generators in terms of geodesic polar and Beltrami (projective) coordinates will enable us to define the curved counterparts of these two systems.

2.1 Geodesic polar coordinates

Let $\mathfrak{so}_\kappa(3)$ be the 3D real Lie algebra with generators J_{01}, J_{02} and J_{12} with commutation relations and Casimir operator given by

$$[J_{12}, J_{01}] = J_{02}, \quad [J_{12}, J_{02}] = -J_{01}, \quad [J_{01}, J_{02}] = \kappa J_{12}, \quad \mathcal{C} = J_{01}^2 + J_{02}^2 + \kappa J_{12}^2, \tag{2.4}$$

where κ is a real parameter. The three 2D classical Riemannian spaces of constant Gaussian curvature κ can be collectively constructed as the homogenous spaces $SO_\kappa(3)/SO(2)$ where $SO_\kappa(3)$ is the Lie group generated by $\mathfrak{so}_\kappa(3)$ and $SO(2)$ is the isotopy subgroup spanned by J_{12}. Hence J_{01} and J_{02} play the role of generators of translations, while J_{12} corresponds to the generator of rotations on the 2D space. In this way, according to the value of κ we get the three particular homogeneous spaces:

$\kappa > 0$: Sphere	$\kappa = 0$: Euclidean plane	$\kappa < 0$: Hyperbolic plane
$\mathbf{S}^2 = SO(3)/SO(2)$	$\mathbf{E}^2 = ISO(2)/SO(2)$	$\mathbf{H}^2 = SO(2,1)/SO(2)$



Figure 1: Schematic representation of the symmetry generators J_{ij} and geodesic distances of a point Q with geodesic polar coordinates (r,ϕ) on a 2D homogeneous space.

We recall that J_{12} leaves a point O invariant, the origin, and that each translation generator J_{0i} moves O along a base geodesic l_i in such a manner that l_1 and l_2 are orthogonal at O as depicted in figure 1.

These spaces can be globally embedded in the linear space $\mathbb{R}^3$ with ambient or Weierstrass coordinates $(x_0,\mathbf{x})=(x_0,x_1,x_2)$ by imposing the "sphere" constraint Σ: $x_0^2+\kappa\,\mathbf{x}^2=1$; the origin corresponds to the point $O=(1,\mathbf{0})\in\mathbb{R}^3$. The geodesic polar coordinates $r\in(0,\infty)$ and $\phi\in[0,2\pi)$ are defined as follows. If the particle is located at a point Q, then r is the distance between Q and the origin O measured along the geodesic l that joins both points. On the other hand, ϕ is the angle which determines the orientation of l with respect to the base geodesic l_1 (see figure 1). The ambient coordinates are parametrized in terms of (r,ϕ) in the form [11, 15]

$$x_0=\mathrm{C}_\kappa(r),\qquad x_1=\mathrm{S}_\kappa(r)\cos\phi,\qquad x_2=\mathrm{S}_\kappa(r)\sin\phi,$$

where we have introduced the curvature-dependent functions

$$\mathrm{C}_\kappa(r)=\left\{\begin{array}{ll}\cos\sqrt{\kappa}\,r & \kappa>0\\ 1 & \kappa=0\\ \cosh\sqrt{-\kappa}\,r & \kappa<0\end{array}\right.,\qquad \mathrm{S}_\kappa(r)=\left\{\begin{array}{ll}\frac{1}{\sqrt{\kappa}}\sin\sqrt{\kappa}\,r & \kappa>0\\ r & \kappa=0\\ \frac{1}{\sqrt{-\kappa}}\sinh\sqrt{-\kappa}\,r & \kappa<0\end{array}\right..$$

The κ-tangent is defined by $\mathrm{T}_\kappa(r)=\mathrm{S}_\kappa(r)/\,\mathrm{C}_\kappa(r)$. Properties for these κ-functions can be found in [11, 15, 16]. For instance,

$$\mathrm{C}^2_\kappa(r)+\kappa\,\mathrm{S}^2_\kappa(r)=1,\qquad \frac{\mathrm{d}}{\mathrm{d}r}\,\mathrm{C}_\kappa(r)=-\kappa\,\mathrm{S}_\kappa(r),\qquad \frac{\mathrm{d}}{\mathrm{d}r}\,\mathrm{S}_\kappa(r)=\,\mathrm{C}_\kappa(r).$$

Now let us denote by (p_r,p_ϕ) the conjugate momenta of (r,ϕ). The canonical Poisson bracket for the three functions

$$J_{01}=\cos\phi\,p_r-\frac{\sin\phi}{\mathrm{T}_\kappa(r)}\,p_\phi,\qquad J_{02}=\sin\phi\,p_r+\frac{\cos\phi}{\mathrm{T}_\kappa(r)}\,p_\phi,\qquad J_{12}=p_\phi \tag{2.5}$$

gives us the Lie–Poisson analogue of the algebra $\mathfrak{so}_\kappa(3)$ (2.4). Therefore, (2.5) provides a symplectic realization of the Lie–Poisson analogue of $\mathfrak{so}_\kappa(3)$, and we can write the kinetic energy $\mathcal{T}$ for the free motion of a particle moving on $\mathbf{S}^2$ and $\mathbf{H}^2$ as the above symplectic realization for the Casimir function of $\mathfrak{so}_\kappa(3)$, namely:

$$\mathcal{T} \equiv \frac{1}{2}\mathcal{C} = \frac{1}{2}(J_{01}^2 + J_{02}^2 + \kappa J_{12}^2) = \frac{1}{2}\left(p_r^2 + \frac{p_\phi^2}{\mathrm{S}_\kappa^2(r)}\right). \tag{2.6}$$

Obviously, the $\kappa \to 0$ limit of this expression gives the 2D Euclidean kinetic energy.

2.2 Beltrami coordinates

On the other hand, if we apply the central projection with pole $(0, \mathbf{0}) \in \mathbb{R}^3$ from $(x_0, \mathbf{x})$ to the Beltrami coordinates $\mathbf{q} \in \mathbb{R}^2$, that are defined as $(0, \mathbf{0}) + \mu\,(1, \mathbf{q}) \in \Sigma$, we get the relations

$$\mu = \frac{1}{\sqrt{1+\kappa\mathbf{q}^2}}, \qquad x_0 = \mu, \qquad \mathbf{x} = \mu\,\mathbf{q} = \frac{\mathbf{q}}{\sqrt{1+\kappa\mathbf{q}^2}}.$$

The image of this projection is the subset of $\mathbb{R}^3$ with $\mu \in \mathbb{R}$ so that $1 + \kappa\mathbf{q}^2 > 0$. In the hyperbolic or Lobachevsky plane with $\kappa = -|\kappa| < 0$ this gives the interior of the Poincaré disk: $1 > |\kappa|\mathbf{q}^2$. Notice that when $\kappa = 0$, the Beltrami coordinates reduce to the Cartesian ones on $\mathbf{E}^2$: $x_0 = 1$ and $\mathbf{x} = \mathbf{q}$. Hereafter we assume the following notation for $\mathbf{q} = (q_1, q_2)$ and for the conjugate momenta $\mathbf{p} = (p_1, p_2)$:

$$\mathbf{q}^2 = q_1^2 + q_2^2, \qquad \mathbf{p}^2 = p_1^2 + p_2^2, \qquad \mathbf{q}\cdot\mathbf{p} = q_1 p_1 + q_2 p_2, \qquad |\mathbf{q}| = \sqrt{q_1^2 + q_2^2}.$$

Then another symplectic realization of the Lie–Poisson algebra $\mathfrak{so}_\kappa(3)$ is found to be [17]

$$J_{0i} = p_i + \kappa(\mathbf{q}\cdot\mathbf{p})q_i, \quad i = 1, 2; \qquad J_{12} = q_1 p_2 - q_2 p_1, \tag{2.7}$$

and the free Hamiltonian $\mathcal{T}$ that determines the geodesic motion on the Riemannian spaces is again obtained from the Casimir (2.4) under the above realization, namely

$$\mathcal{T} \equiv \frac{1}{2}\mathcal{C} = \frac{1}{2}(J_{01}^2 + J_{02}^2 + \kappa J_{12}^2) = \frac{1}{2}\left(1 + \kappa\mathbf{q}^2\right)\left(\mathbf{p}^2 + \kappa(\mathbf{q}\cdot\mathbf{p})^2\right). \tag{2.8}$$

We remark that, as expected, there is canonical equivalence between the Beltrami and geodesic polar variables through the canonical transformation given by

$$\begin{aligned} q_1 &= \mathrm{T}_\kappa(r)\cos\phi, \qquad & p_1 &= \mathrm{C}_\kappa^2(r)\cos\phi\, p_r - \frac{\sin\phi}{\mathrm{T}_\kappa(r)}\, p_\phi, \\ q_2 &= \mathrm{T}_\kappa(r)\sin\phi, \qquad & p_2 &= \mathrm{C}_\kappa^2(r)\sin\phi\, p_r + \frac{\cos\phi}{\mathrm{T}_\kappa(r)}\, p_\phi, \end{aligned} \tag{2.9}$$

which generalizes (2.1). The inverse change of coordinates leads to the expressions

$$\begin{aligned} &\mathrm{C}_\kappa(r) = \frac{1}{\sqrt{1+\kappa\mathbf{q}^2}}, \qquad && \mathrm{S}_\kappa(r) = \frac{|\mathbf{q}|}{\sqrt{1+\kappa\mathbf{q}^2}}, \qquad && \mathrm{T}_\kappa(r) = |\mathbf{q}|, \\ &\cos\phi = \frac{q_1}{|\mathbf{q}|}, && \sin\phi = \frac{q_2}{|\mathbf{q}|}, && \tan\phi = \frac{q_2}{q_1}. \end{aligned} \tag{2.10}$$

3 The anisotropic Higgs oscillator

In order to generalize the integrable anisotropic Hamiltonian on $\mathbf{E}^2$ (2.2) to the sphere and the hyperbolic plane we recall that, in terms of geodesic polar coordinates, the Higgs oscillator potential on both spaces is given by $\mathrm{T}_\kappa^2(r)$ [11, 12, 13] (i.e. $\tan^2 r$ on $\mathbf{S}^2$ with $\kappa = 1$ and $\tanh^2 r$ on $\mathbf{H}^2$ with $\kappa = -1$). Furthermore, the superposition of the Higgs oscillator with two centrifugal potentials (i.e. the curved SW system) is also known in these variables and defines a superintegrable Hamiltonian. Such a system will be our starting point, whose properties are summarized as follows.

Proposition 1. [11, 12, 13] *The Higgs oscillator Hamiltonian on* $\mathbf{S}^2$ *and* $\mathbf{H}^2$ *with two Rosochatius terms (curved SW system) is given in terms of geodesic polar variables by*

$$\mathcal{H}_{\mathrm{SW}} = \frac{1}{2}\left(p_r^2 + \frac{p_\phi^2}{\mathrm{S}_\kappa^2(r)}\right) + \delta\,\mathrm{T}_\kappa^2(r) + \frac{\lambda_1}{\mathrm{S}_\kappa^2(r)\cos^2\phi} + \frac{\lambda_2}{\mathrm{S}_\kappa^2(r)\sin^2\phi}. \tag{3.1}$$

The Hamiltonian $\mathcal{H}_{\mathrm{SW}}$ *Poisson-commutes with the functions*

$$\begin{aligned} I_{01} &= \frac{1}{2}J_{01}^2 + \delta\,\mathrm{T}_\kappa^2(r)\cos^2\phi + \frac{\lambda_1}{\mathrm{T}_\kappa^2(r)\cos^2\phi}, \\ I_{02} &= \frac{1}{2}J_{02}^2 + \delta\,\mathrm{T}_\kappa^2(r)\sin^2\phi + \frac{\lambda_2}{\mathrm{T}_\kappa^2(r)\sin^2\phi}, \\ I_{12} &= \frac{1}{2}J_{12}^2 + \lambda_1\tan^2\phi + \frac{\lambda_2}{\tan^2\phi}, \end{aligned} \tag{3.2}$$

where J_{ij} *are the generators (2.5). Each set* $(\mathcal{H}_{\mathrm{SW}}, I_{01}, I_{12})$ *and* $(\mathcal{H}_{\mathrm{SW}}, I_{02}, I_{12})$ *is formed by three functionally independent functions. Moreover,* $\mathcal{H}_{\mathrm{SW}} = I_{01} + I_{02} + \kappa I_{12} + \kappa(\lambda_1 + \lambda_2)$.

Essentially, our task now consists in the introduction of an anharmonic Ω-term in the Hamiltonian (3.1) by keeping the integrability of the system. This process requires to modify simultaneously the integrals of motion (3.2). The final result is presented in the following proposition, that can be proven through direct computations.

Proposition 2. *The anisotropic Higgs oscillator Hamiltonian on* $\mathbf{S}^2$ *and* $\mathbf{H}^2$ *with two Rosochatius terms defined in geodesic polar variables as*

$$\mathcal{H}_\kappa = \frac{1}{2}\left(p_r^2 + \frac{p_\phi^2}{\mathrm{S}_\kappa^2(r)}\right) + \delta\,\mathrm{T}_\kappa^2(r) + \Omega\,\mathrm{S}_\kappa^2(r)\sin^2\phi + \frac{\lambda_1}{\mathrm{S}_\kappa^2(r)\cos^2\phi} + \frac{\lambda_2}{\mathrm{S}_\kappa^2(r)\sin^2\phi} \tag{3.3}$$

Poisson-commutes with the functions

$$\begin{aligned} \mathcal{I}_{1,\kappa} &= \frac{1}{2}J_{01}^2 + \delta\,\mathrm{T}_\kappa^2(r)\cos^2\phi + \frac{\lambda_1}{\mathrm{T}_\kappa^2(r)\cos^2\phi}, \\ \mathcal{I}_{2,\kappa} &= \frac{1}{2}\left(J_{02}^2 + \kappa J_{12}^2\right) + \delta\,\mathrm{T}_\kappa^2(r)\sin^2\phi + \Omega\,\mathrm{S}_\kappa^2(r)\sin^2\phi + \frac{\lambda_2}{\mathrm{T}_\kappa^2(r)\sin^2\phi} \\ &\qquad + \kappa\left(\frac{\lambda_1}{\cos^2\phi} + \frac{\lambda_2}{\sin^2\phi}\right), \end{aligned} \tag{3.4}$$

where J_{ij} *are the generators (2.5). The pairs* $(\mathcal{H}_\kappa, \mathcal{I}_{1,\kappa})$ *and* $(\mathcal{H}_\kappa, \mathcal{I}_{2,\kappa})$ *are constituted by two functionally independent functions, and* $\mathcal{H}_\kappa = \mathcal{I}_{1,\kappa} + \mathcal{I}_{2,\kappa}$.



(a) (b) (c) (d)

Figure 2: Some trajectories of the Hamiltonians $\mathcal{H}_\kappa$ (3.3) and $\mathcal{H}'_\kappa$ (4.1) on the sphere $\mathbf{S}^2$ with $\kappa = 1$ and without centrifugal terms ($\lambda_1 = \lambda_2 = 0$). They are plotted in $\mathbb{R}^3$ with ambient coordinates $(x_0, \mathbf{x})$ such that $x_0^2 + x_1^2 + x_2^2 = 1$. Time runs from $t = 0$ to $t = 8$ for the initial data $q_1 = 1$, $\dot{q}_1 = 1$, $q_2 = -0.5$, $\dot{q}_2 = 2$. (a) the superintegrable case of $\mathcal{H}_\kappa$ with $\delta = 1$ and $\Omega = 0$ (Higgs oscillator); (b) the integrable $\mathcal{H}_\kappa$ with $\delta = 1$ and $\Omega = 0.2$; (c) the integrable $\mathcal{H}_\kappa$ with $\delta = 1$ and $\Omega = 3$; and (d) the superintegrable $\mathcal{H}'_\kappa$ with $\delta = 1$.

We stress that for a generic value of Ω the anisotropic Higgs oscillator $\mathcal{H}_\kappa$ (3.3) is an integrable (but, presumably, non superintegrable) Ω-anharmonic generalization of $\mathcal{H}_{\mathrm{SW}}$ (3.1) since the former is endowed with a single quadratic integral of motion (either $\mathcal{I}_{1,\kappa}$ or $\mathcal{I}_{2,\kappa}$) and its flat $\kappa \to 0$ limit is not superintegrable. We remark that in this generalization only $\mathcal{I}_{1,\kappa} \equiv I_{01}$ "survives" from the set of three integrals (3.2), meanwhile the two remaining ones are mixed within $\mathcal{I}_{2,\kappa}$, and when $\Omega = 0$ we get $\mathcal{I}_{2,\kappa} = I_{02} + \kappa I_{12} + \kappa(\lambda_1 + \lambda_2)$. Note also that the $\kappa = 0$ Euclidean limit is naturally included in (3.3) and (3.4), and provides (2.2).

The system defined by proposition 2 can also be written in terms of Beltrami variables by applying the transformation (2.10), and it contains only polynomial and rational expresions

$$\begin{aligned}
\mathcal{H}_\kappa &= \frac{1}{2}\left(1 + \kappa\mathbf{q}^2\right)\left(\mathbf{p}^2 + \kappa(\mathbf{q}\cdot\mathbf{p})^2\right) + \delta\mathbf{q}^2 + \Omega\,\frac{q_2^2}{(1+\kappa\mathbf{q}^2)} + \left(1 + \kappa\mathbf{q}^2\right)\left(\frac{\lambda_1}{q_1^2} + \frac{\lambda_2}{q_2^2}\right), \\
\mathcal{I}_{1,\kappa} &= \frac{1}{2}J_{01}^2 + \delta q_1^2 + \frac{\lambda_1}{q_1^2}, \qquad\qquad (3.5) \\
\mathcal{I}_{2,\kappa} &= \frac{1}{2}\left(J_{02}^2 + \kappa J_{12}^2\right) + \delta q_2^2 + \Omega\,\frac{q_2^2}{(1+\kappa\mathbf{q}^2)} + \frac{\lambda_2}{q_2^2} + \kappa\mathbf{q}^2\left(\frac{\lambda_1}{q_1^2} + \frac{\lambda_2}{q_2^2}\right),
\end{aligned}$$



(a)

(b)

(c)

(d)

Figure 3: Some trajectories of the Hamiltonians $\mathcal{H}_\kappa$ (3.3) and $\mathcal{H}'_\kappa$ (4.1) on the hyperboloid $\mathbf{H}^2$ with $\kappa = -1$ and without centrifugal terms ($\lambda_1 = \lambda_2 = 0$). They are plotted in $\mathbb{R}^3$ with ambient coordinates $(x_0, \mathbf{x})$ such that $x_0^2 - x_1^2 - x_2^2 = 1$. Time runs from $t = 0$ to $t = 12$ with the initial data $q_1 = 0.1$, $\dot{q}_1 = 0.1$, $q_2 = 0.3$, $\dot{q}_2 = -0.2$. (a) the superintegrable case of $\mathcal{H}_\kappa$ with $\delta = 1$ and $\Omega = 0$ (Higgs oscillator); (b) the integrable $\mathcal{H}_\kappa$ with $\delta = 1$ and $\Omega = 0.2$; (c) the integrable $\mathcal{H}_\kappa$ with $\delta = 1$ and $\Omega = 3$; and (d) the superintegrable $\mathcal{H}'_\kappa$ with $\delta = 1$.

where now J_{ij} are the functions given by (2.7). In this way, the curved Higgs oscillator potential, $\mathrm{T}^2_\kappa(r)$, becomes the simple "flat" expression $\mathbf{q}^2$ and the Euclidean Hamiltonian $\mathcal{H}$ is recovered in Cartesian coordinates, in the form (1.1) and (1.2), by setting $\kappa = 0$ in (3.5).

Numerical solutions for some particular bounded trajectories of $\mathcal{H}_\kappa$ (3.3) are explicitly illustrated in figures 2 (2D sphere) and 3 (hyperboloid). They are obtained by fixing the values of the parameters Ω and δ (for the sake of simplicity both λ_i are set equal to zero) and by taking the same initial conditions. In this respect, it is worth remarking that:

- The main dynamical footprint of superintegrable systems consists in the fact that all their bounded trajectories are periodic ones. Indeed, when $\Omega = \lambda_1 = \lambda_2 = 0$ we recover the superintegrable Higgs oscillator, whose periodic trajectories are plotted in figures 2(a) and 3(a).

- The integrable Hamiltonian (3.3), arising by introducing the anharmonic potential with $\Omega \neq 0$ (we keep $\lambda_1 = \lambda_2 = 0$), corresponds to the anisotropic Higgs oscillator. In this case the bounded trajectories shown in figures 2(b) and 3(b) turn out to be nonperiodic. Therefore, the transition from the Higgs oscillator ($\Omega = 0$) to the anisotropic one

($\Omega \neq 0$) can be graphically appreciated by comparing figures 2(a) and 2(b) (see also 3(a) and 3(b)).

- We also analyse the particular case of $\mathcal{H}_\kappa$ with $\Omega = 3\delta$ and $\lambda_1 = \lambda_2 = 0$, which is a curved analogue ($\kappa \neq 0$) of the superintegrable Euclidean oscillator 1:2 ($\kappa = 0$) whose flat trajectory is the Lissajous 1:2 curve. For the initial conditions here chosen, it is found that the curved bounded trajectories are no longer periodic as it is depicted in figures 2(c) and 3(c).

Therefore, the Hamiltonian $\mathcal{H}_\kappa$ (3.3) seems to be superintegrable only in the $\Omega = 0$ case, although we stress that we have not rigurously proven the non-superintegrability of $\mathcal{H}_\kappa$ for an arbitrary value of Ω. Equivalently, we could say that $\mathcal{H}_\kappa$ presents a superintegrability breaking induced by the non-vanishing curvature of the underlying space.

4 A superintegrable anisotropic oscillator

As far as the known superintegrable anisotropic system (2.3) is concerned, its curved counterpart for any value of the curvature κ can be written as follows.

Proposition 3. [11] *The Hamiltonian*

$$\begin{aligned} \mathcal{H}'_\kappa = {} & \frac{1}{2}\left(p_r^2 + \frac{p_\phi^2}{\mathrm{S}_\kappa^2(r)}\right) + \delta\, \frac{\mathrm{S}_\kappa^2(r)\cos^2\phi}{\left(1 - \kappa\, \mathrm{S}_\kappa^2(r)\cos^2\phi\right)} \\ & + 4\delta\, \frac{\mathrm{T}_\kappa^2(r)\sin^2\phi}{\left(1 - \kappa\, \mathrm{S}_\kappa^2(r)\cos^2\phi\right)\left(1 - \kappa\, \mathrm{T}_\kappa^2(r)\sin^2\phi\right)^2} + \frac{\lambda_1}{\mathrm{S}_\kappa^2(r)\cos^2\phi} \end{aligned} \tag{4.1}$$

is endowed with three integrals of motion given by

$$\begin{aligned} \mathcal{I}'_{1,\kappa} &= \frac{1}{2}\left(J_{01}^2 + \kappa J_{12}^2\right) + \delta\, \frac{\mathrm{T}_\kappa^2(r)\cos^2\phi\left(1 + \kappa\, \mathrm{T}_\kappa^2(r)\sin^2\phi\right)}{\left(1 - \kappa\, \mathrm{T}_\kappa^2(r)\sin^2\phi\right)^2} + \frac{\lambda_1}{\mathrm{S}_\kappa^2(r)\cos^2\phi}, \\ \mathcal{I}'_{2,\kappa} &= \frac{1}{2}J_{02}^2 + 4\delta\, \frac{\mathrm{T}_\kappa^2(r)\sin^2\phi}{\left(1 - \kappa\, \mathrm{T}_\kappa^2(r)\sin^2\phi\right)^2}, \\ \mathcal{L}'_\kappa &= J_{01}J_{12} + 2\delta\, \frac{\mathrm{T}_\kappa^3(r)\cos^2\phi\sin\phi}{\left(1 - \kappa\, \mathrm{T}_\kappa^2(r)\sin^2\phi\right)^2} - 2\lambda_1\, \frac{\tan\phi}{\mathrm{T}_\kappa(r)\cos\phi}, \end{aligned} \tag{4.2}$$

where J_{ij} are the generators (2.5) and $\mathcal{H}'_\kappa = \mathcal{I}'_{1,\kappa} + \mathcal{I}'_{2,\kappa}$. Both sets $(\mathcal{H}'_\kappa, \mathcal{I}'_{1,\kappa}, \mathcal{L}'_\kappa)$ and $(\mathcal{H}'_\kappa, \mathcal{I}'_{2,\kappa}, \mathcal{L}'_\kappa)$ are formed by three functionally independent functions.

Once again, this system can be more easily expressed by introducing Beltrami variables through (2.10). In this way we find that

$$\begin{aligned} \mathcal{H}'_\kappa &= \frac{1}{2}\left(1 + \kappa\mathbf{q}^2\right)\left(\mathbf{p}^2 + \kappa(\mathbf{q}\cdot\mathbf{p})^2\right) + \frac{\delta\, q_1^2}{(1 + \kappa q_2^2)} + \frac{4\delta\,(1 + \kappa\mathbf{q}^2)q_2^2}{(1 + \kappa q_2^2)(1 - \kappa q_2^2)^2} + \lambda_1\, \frac{(1 + \kappa\mathbf{q}^2)}{q_1^2}, \\ \mathcal{I}'_{1,\kappa} &= \frac{1}{2}\left(J_{01}^2 + \kappa J_{12}^2\right) + \delta\, \frac{q_1^2(1 + \kappa q_2^2)}{(1 - \kappa q_2^2)^2} + \lambda_1\, \frac{(1 + \kappa\mathbf{q}^2)}{q_1^2}, \\ \mathcal{I}'_{2,\kappa} &= \frac{1}{2}J_{02}^2 + 4\delta\, \frac{q_2^2}{(1 - \kappa q_2^2)^2}, \qquad \mathcal{L}'_\kappa = J_{01}J_{12} + 2\delta\, \frac{q_1^2 q_2}{(1 - \kappa q_2^2)^2} - 2\lambda_1\, \frac{q_2}{q_1^2}, \end{aligned} \tag{4.3}$$



(a) (b)

(c) (d)

Figure 4: The potentials $\mathcal{U}$ (3.5) and $\mathcal{U}'$ (4.3) (without centrifugal terms $\lambda_1 = \lambda_2 = 0$) in Beltrami coordinates on the sphere $\mathbf{S}^2$ with $\kappa = 1$ and on the hyperbolic plane $\mathbf{H}^2$ with $\kappa = -1$. (a) the integrable $\mathcal{U}$ on $\mathbf{S}^2$ for $\delta = 1$ and $\Omega = 3$; (b) the superintegrable $\mathcal{U}'$ on $\mathbf{S}^2$ for $\delta = 1$; (c) the integrable $\mathcal{U}$ on $\mathbf{H}^2$ for $\delta = 1$ and $\Omega = 3$; and (d) the superintegrable $\mathcal{U}'$ on $\mathbf{H}^2$ for $\delta = 1$.

where J_{ij} are now the generators (2.7). In this variables, the Euclidean case with $\kappa = 0$ leads just to the equations (1.3) defining $\mathcal{H}'$ in Cartesian coordinates.

The numerical integration of the trajectories for the superintegrable Hamiltonian $\mathcal{H}'_\kappa$ (4.1) gives rise to the Lissajous 1:2 curves on the sphere and the hyperboloid which are shown in figures 2(d) and 3(d), respectively. They are worthy to be compared with figures 2(c) and 3(c). Hence, in contrast with the integrable Hamiltonian $\mathcal{H}_\kappa$ (3.3) with $\Omega = 3\delta$, the system $\mathcal{H}'_\kappa$ can be regarded as the appropriate curved analogue ($\kappa \neq 0$) of the anisotropic Euclidean oscillator 1:2 ($\kappa = 0$). We recall that in the Euclidean case $\mathcal{H}'$ (1.3) is just the particular case of $\mathcal{H}$ (1.1) with $\Omega = 3\delta$ and $\lambda_2 = 0$, but this is no longer true when the curvature is turned on. Consequently, we have shown that the Euclidean 1:2 oscillator admits (at least) two different integrable curved generalizations, since $\mathcal{H}'_\kappa$ (the superintegrable one) cannot be recovered as a particular case of $\mathcal{H}_\kappa$ with $\Omega = 3\delta$ and $\lambda_2 = 0$.

The difference between these two systems can also be highlighted by representing the corresponding potentials. We write $\mathcal{H}_\kappa = \mathcal{T} + \mathcal{U}$ (3.5) and $\mathcal{H}'_\kappa = \mathcal{T} + \mathcal{U}'$ (4.3), with $\mathcal{T}$ given by (2.8), and we represent both potentials $\mathcal{U}$ (with $\Omega = 3\delta$) and $\mathcal{U}'$ in Beltrami coordinates in figure 4. Notice that the corresponding trajectories for the potentials shown in figures 4(a), 4(b), 4(c) and 4(d) are drawn in figures 2(c), 2(d), 3(c) and 3(d), respectively.

On the other hand, the differences between these two curved anisotropic potentials can also be enhanced by rewritting them in terms of the ambient coordinates $(x_0, \mathbf{x})$ introduced in section 2, namely

$$\begin{aligned} \mathcal{U} &= \delta \, \frac{\mathbf{x}^2}{(1-\kappa \mathbf{x}^2)} + \Omega \, x_2^2 + \frac{\lambda_1}{x_1^2} + \frac{\lambda_2}{x_2^2}, \\ \mathcal{U}' &= \delta \, \frac{x_1^2}{(1-\kappa x_1^2)} + 4\delta \, \frac{x_0^2 x_2^2}{(x_0^2+\kappa x_2^2)(x_0^2-\kappa x_2^2)^2} + \frac{\lambda_1}{x_1^2}, \end{aligned} \tag{4.4}$$

and recall that $x_0 \to 1$ when $\kappa \to 0$. In this form, these potentials can be more easily compared with the proposals for integrable anharmonic oscillators on the spheres and hyperbolic spaces that have been previously presented in [18, 19], thus concluding that the anisotropic Higgs oscillator potential $\mathcal{U}$ defines a new integrable system. In fact, the potentials studied in [18, 19] are just polynomials when written in ambient coordinates. For instance, the integrable potential on the n-dimensional sphere considered in [19] reads

$$\mathcal{V} = \sum_{i=0}^{n} a_i^2 x_i^2 - \left(\sum_{i=0}^{n} a_i x_i^2 \right)^2 ,$$

where a_i are arbitrary real constants and x_i are $(n+1)$ ambient coordinates such that $\sum_{i=0}^{n} x_i^2 = 1$. The systems worked out in [18] have a similar form. Evidently, they are quite different from (4.4).

5 Curved "centrifugal" terms: geometry and dynamics

It is worth recalling that the SW system on $\mathbf{S}^2$ was interpreted in [20, 21, 22] as a superposition of three spherical oscillators, and this result has been extended to higher dimensions and to other spaces with constant curvature in [12, 13, 23]. The crucial point is to take into account that the Higgs oscillator potential given by $\mathrm{T}_\kappa^2(r)$, is a *central* harmonic oscillator, whose center is located at the origin O, and r is the distance between Q and O measured along the geodesic l that joins both points (see section 2). Hence if, instead of O, we take a generic point Q_ρ such that ρ is the distance between Q and Q_ρ measured along the geodesic l_ρ that joins both points, then $\mathrm{T}_\kappa^2(\rho)$ can be interpreted as a *noncentral* harmonic oscillator potential with center at Q_ρ. With this ideas in mind, let us consider the potentials $\mathcal{U}$ and $\mathcal{U}'$ written in geodesic polar coordinates (3.3) and (4.1) in order to search terms of the type $\mathrm{T}_\kappa^2(\rho)$.

Consider the two base geodesics l_1 and l_2 orthogonal at the origin O and the point $Q(r, \phi)$. Let Q_1 (resp. Q_2) be the intersection point of l_2 (resp. l_1) with its orthogonal geodesic l_1' (resp. l_2'), $x = QQ_1$ the distance measured along l_1' and $y = QQ_2$ the one measured along l_2' as depicted in figure 1. Next we apply the sine theorem [16] on the orthogonal triangles OQQ_1 (with external angle ϕ) and OQQ_2 (with inner angle ϕ) which yields

$$OQQ_1: \quad \mathrm{S}_\kappa(x) = \mathrm{S}_\kappa(r)\cos\phi\,; \qquad OQQ_2: \quad \mathrm{S}_\kappa(y) = \mathrm{S}_\kappa(r)\sin\phi.$$

Therefore the integrable potential $\mathcal{U}$ (3.3) can be expressed as

$$\mathcal{U} = \delta\, \mathrm{T}_\kappa^2(r) + \Omega\, \mathrm{S}_\kappa^2(y) + \frac{\lambda_1}{\mathrm{S}_\kappa^2(x)} + \frac{\lambda_2}{\mathrm{S}_\kappa^2(y)}, \tag{5.1}$$

which provides a unified description of this potential on the three Riemannian spaces:

- The δ-term is a central harmonic oscillator with center at the origin O.
- The Ω-term is the anharmonic potential which only depends on the geodesic distance y to the point Q_2, which is not a fixed center since it is moving together with the particle located at Q.
- The λ_i-terms ($i = 1, 2$) are two “centrifugal barriers”.

Likewise, the superintegrable potential $\mathcal{U}'$ (4.1) can be rewritten as

$$\mathcal{U}' = \delta\, \mathrm{T}^2_\kappa(x) + 4\delta\, \mathrm{T}^2_\kappa(y) \left(\frac{\mathrm{C}^2_\kappa(r)\, \mathrm{C}^2_\kappa(y)}{\mathrm{C}^2_\kappa(x) \left(\mathrm{C}^2_\kappa(r) - \kappa\, \mathrm{S}^2_\kappa(y) \right)^2} \right) + \frac{\lambda_1}{\mathrm{S}^2_\kappa(x)}, \tag{5.2}$$

showing that this potential is not a simple addition of some anharmonic potential (only depending on the variable y) to the *central* harmonic oscillator; its interpretation is as follows:

- The first term is a noncentral harmonic oscillator with center at the point Q_1 (which is, moreover, not fixed).
- The second one is not any kind of curved harmonic oscillator, although it could be regarded as some kind of “deformation” of a noncentral oscillator with center at Q_2.
- The λ_1-term is a “centrifugal barrier”.

It is worth stressing that the Euclidean limit is well defined in both (5.1) and (5.2), namely

$$\kappa \to 0: \qquad \mathcal{U} = \delta r^2 + \Omega y^2 + \frac{\lambda_1}{x^2} + \frac{\lambda_2}{y^2}, \qquad \mathcal{U}' = \delta x^2 + 4\delta y^2 + \frac{\lambda_1}{x^2}, \tag{5.3}$$

where now the geodesic distances x and y become Cartesian coordinates (that is, q_1 and q_2 appearing in expressions (1.1) and (1.3)).

Furthermore, on the sphere $\mathbf{S}^2$ the potential $\mathcal{U}$ admits a second interpretation, which requires to rewrite the “centrifugal barriers” as noncentral harmonic oscillators. Explicitly, let us consider the potential $\mathcal{U}$ in the form (5.1), set $\kappa = +1$ for simplicity and let O_i be the *fixed* point placed along the geodesic l_i ($i = 1, 2$) which is at a distance $\frac{\pi}{2}$ from the origin O. Notice that in ambient coordinates $(x_0, \mathbf{x})$ we have that $O = (1, 0, 0)$, $O_1 = (0, 1, 0)$ and $O_2 = (0, 0, 1)$. We introduce the distance r_i between Q and O_i measured along l'_i such that the three points $\{O_i Q Q_i\}$ lie on the same geodesic l'_i as it is shown in figure 5(a) in the ambient space $\mathbb{R}^3$. Thus we obtain that

$$r_1 + x = r_2 + y = \frac{\pi}{2}, \qquad x_0 = \cos r, \qquad x_1 = \sin x = \cos r_1, \qquad x_2 = \sin y = \cos r_2.$$

Then the potential (5.1) on $\mathbf{S}^2$ can be expressed as

$$\begin{aligned} \mathcal{U} &= \delta \tan^2 r + \Omega \sin^2 y + \frac{\lambda_1}{\sin^2 x} + \frac{\lambda_2}{\sin^2 y} \\ &= \delta \tan^2 r + \frac{\Omega}{(1 + \tan^2 r_2)} + \lambda_1 \tan^2 r_1 + \lambda_2 \tan^2 r_2 + \lambda_1 + \lambda_2. \end{aligned} \tag{5.4}$$

Hence this corresponds to the superposition of the central (Higgs) oscillator with center at O with an anharmonic potential depending on the distance r_2 to O_2 and with two more



(a) (b)

Figure 5: Geodesic distances involved in the geometric description of the integrable anisotropic Higgs oscillator with two Rosochatius terms $\mathcal{U}$: (a) on the sphere for the expression (5.4) and (b) on the hyperboloid in the form (5.5).

noncentral oscillators determined by the geodesic distances r_i from the centers placed at the fixed points O_i. Notice that the flat limit $\kappa = 0$ can only be performed in the former expression of (5.4) providing (5.3) with the two Euclidean centrifugal barriers, since in the latter the distances $r_i \to \infty$ (that is, the centers O_i go to infinity) so that the terms $\tan^2 r_i$ become indeterminate.

We remark that in the hyperbolic plane with $\kappa < 0$ a similar description in terms of oscillators is, in principle, precluded. The analogous points to the previous centers O_i would be beyond infinity as it can be seen in figure 5(b), that is, beyond the "proper" hyperbolic space $\mathbf{H}^2$. In particular, if we set $\kappa = -1$ we find that

$$x_0 = \cosh r, \qquad x_1 = \sinh x, \qquad x_2 = \sinh y,$$

and the potential (5.1) on $\mathbf{H}^2$ reads as

$$\begin{aligned} \mathcal{U} &= \delta \tanh^2 r + \Omega \sinh^2 y + \frac{\lambda_1}{\sinh^2 x} + \frac{\lambda_2}{\sinh^2 y} \\ &= \delta \tanh^2 r + \Omega \sinh^2 y + \frac{\lambda_1}{\tanh^2 x} + \frac{\lambda_2}{\tanh^2 y} + \lambda_1 + \lambda_2. \end{aligned} \tag{5.5}$$

Hence the λ_i-terms only admit an interpretation as centrifugal barriers with Euclidean limit given by (5.3). Nevertheless, an interpretation of the λ_i-potentials as noncentral oscillators can be established if one considers the *complete* hyperbolic space. In that case, the centers O_i would be located in the "ideal" (exterior) region of $\mathbf{H}^2$ so beyond the "actual" $\mathbf{H}^2$.

5.1 Some trajectories

In order to illustrate the above results and to highlight the role of the curved centrifugal terms, we plot in figure 6 some trajectories of the Hamiltonians $\mathcal{H}_\kappa$ (3.3) and $\mathcal{H}'_\kappa$ (4.1) on the sphere $\mathbf{S}^2$ in ambient coordinates and with a single centrifugal term ($\lambda_1 > 0$ and $\lambda_2 = 0$). These trajectories are numerically integrated by imposing the same initial conditions and values of the constants δ and Ω as in the trajectories plotted in figure 2 (which is the same system with no centrifugal term). Consequently, the effect of the λ_1-potential becomes apparent by



Figure 6: Some trajectories for the Hamiltonians $\mathcal{H}_\kappa$ (3.3) and $\mathcal{H}'_\kappa$ (4.1) on $\mathbf{S}^2$ ($\kappa = 1$) with a single Rosochatius potential such that $\lambda_1 = 0.1$ and $\lambda_2 = 0$. Time runs from $t = 0$ to $t = 8$ with the initial data $q_1 = 1$, $\dot{q}_1 = 1$, $q_2 = -0.5$, $\dot{q}_2 = 2$. (a) the superintegrable case of $\mathcal{H}_\kappa$ with $\delta = 1$ and $\Omega = 0$ (Higgs oscillator with one centrifugal term); (b) the integrable $\mathcal{H}_\kappa$ with $\delta = 1$ and $\Omega = 0.2$; (c) the integrable $\mathcal{H}_\kappa$ with $\delta = 1$ and $\Omega = 3$; and (d) the superintegrable $\mathcal{H}'_\kappa$ with $\delta = 1$.

comparing figures 2 and 6. In this respect, it has to be taken into account that the λ_1-term is associated to the ambient coordinate x_1 (see (4.4)), with center $O_1 = (0, 1, 0)$ for the noncentral oscillator and geodesic radial distance r_1. In this way, we find that:

- The trajectories are restricted to the "right" semisphere due to the infinite barrier arising at $x_1 = 0$. Roughly speaking, this barrier induces a "folding" of the initial trajectories in figure 2.
- The superintegrable Higgs oscillator ($\Omega = 0$) with a centrifugal term provides a kind of Lissajous 1:2 curve (figure 6(a)).
- The integrable anisotropic systems $\mathcal{H}_\kappa$ with $\Omega \neq 0$ represented in figures 6(b) and 6(c) show bounded trajectories which are again nonperiodic.
- The superintegrable $\mathcal{H}'_\kappa$ with a centrifugal term/noncentral oscillator gives rise to a kind of Lissajous 1:1 curve (figure 6(d)).

Next, if the second centrifugal term with $\lambda_2 > 0$ is also considered, then similar features arise from the second infinite barrier. For the sake of completeness, figure 7 contains the



(a) (b) (c)

Figure 7: Trajectories of the Hamiltonian $\mathcal{H}_\kappa$ (3.3) on $\mathbf{S}^2$ ($\kappa = 1$) with two Rosochatius potentials given by $\lambda_1 = 0.1$ and $\lambda_2 = 0.05$. Initial data are the same as in figures 2 and 6. (a) the superintegrable case of $\mathcal{H}_\kappa$ with $\delta = 1$ and $\Omega = 0$ (Higgs oscillator with two centrifugal terms or curved SW system); (b) the integrable $\mathcal{H}_\kappa$ with $\delta = 1$ and $\Omega = 0.2$; and (c) the integrable $\mathcal{H}_\kappa$ with $\delta = 1$ and $\Omega = 3$. Note that $\mathcal{H}'_\kappa$ is not defined when both λ_1 and λ_2 are non-zero.

trajectories on the sphere of the Hamiltonian $\mathcal{H}_\kappa$ (3.3) which exactly correspond to those plotted in figure 6 but now with the second Rosochatius potential. We remark that:

- The trajectories of $\mathcal{H}_\kappa$ are restricted to a quadrant of the sphere; these can be seen as once "folded" with respect to the trajectories shown in figure 6 but twice with respect to those obtained in figure 2.
- The superintegrable Higgs oscillator ($\Omega = 0$) with two centrifugal terms (curved SW system) gives rise to a Lissajous 1:1 curve (figure 7(a)).
- As expected once more, the trajectories coming from the integrable anisotropic Hamiltonian $\mathcal{H}_\kappa$ with $\Omega \neq 0$ drawn in figures 7(b) and 7(c) show bounded nonperiodic trajectories.

Summarizing, we can say that the addition of the appropriate curved centrifugal terms preserves the essential dynamical features of all these systems.

6 Concluding remarks

In this work we have introduced a new integrable Hamiltonian $\mathcal{H}_\kappa$ on $\mathbf{S}^2$ and $\mathbf{H}^2$, that can be interpreted as the superposition of the anisotropic Higgs oscillator with two curved Rosochatius potentials. Furthermore, such a system together with the previously known curved superintegrable Hamiltonian $\mathcal{H}'_\kappa$ have been algebraically and geometrically studied. Numerical solutions for the trajectories corresponding to some particular initial conditions for both Hamiltonians have been obtained. In this respect, some remarks are in order.

- The flat contraction $\kappa \to 0$ is always a well defined and smooth limit in all the expressions presented in the paper. In this way the known systems presented in section 1 can always be recovered in the Euclidean limit. Hence the two different integrable Hamiltonians $\mathcal{H}_\kappa$ and $\mathcal{H}'_\kappa$ can be thought as generalizations of the known Euclidean systems (1.1) and (1.3) (or (2.2) and (2.3)), and $\mathcal{H}'_{\kappa=0}$ is nothing but a particular (superintegrable) case of $\mathcal{H}_{\kappa=0}$.
- Alternatively, this approach can be seen as a "curvature-deformation" process of the Euclidean anharmonic oscillator that by starting from (2.2) and (2.3) arrives at the Hamiltonians (3.3) and (4.1), respectively. In this sense, we remark that there is no an apparent straightforward prescription in order to perform such a deformation, neither in the integrable nor in the superintegrable system, as the explicit expressions for the curved Hamiltonians show.
- It is also worth stressing that when $\kappa \neq 0$ the superintegrable Hamiltonian (4.1) *cannot* be recovered from the integrable one (3.3) by setting $\Omega = 3\delta$ and $\lambda_2 = 0$ (see also the potentials (4.4) as well as the integrals $\mathcal{I}_{i,\kappa}$ (3.4) and $\mathcal{I}'_{i,\kappa}$ (4.3)). This, in turn, means that the Hamiltonians $\mathcal{H}_\kappa$ and $\mathcal{H}'_\kappa$ are two *completely different* systems on the sphere and on the hyperbolic plane. This fact indicates that different integrable curved systems may collapse under contraction into a single flat one and, conversely, that the integrable generalization of a known Euclidean Hamiltonian could split into several generalized "curved" possibilites.

Starting from all these results, several further research directions can be explored. Firstly, the search for superintegrable curved generalizations of other commensurate Euclidean oscillators could be faced. In this respect, we recall that the superintegrable Hamiltonian $\mathcal{H}'$ (1.3) is the only system in the series of anisotropic oscillators with commensurate frequencies that is endowed with quadratic integrals of motion (see [6]). Moreover, the "additional" integral with ensures superintegrability is, in fact, a second order reduction of a true six-th order integral. More explicitly, if we set $\lambda_1 = 0$ then it is known [6] that

$$Q_1 = \left(p_1^2 - 2\delta q_1^2 - 2\mathrm{i}\sqrt{2\delta} q_1 p_1\right)^2 \left(p_2^2 - 8\delta q_2^2 + 4\mathrm{i}\sqrt{2\delta} q_2 p_2\right), \qquad \bar{Q}_1, \qquad Q = \frac{1}{2}\left(Q_1 + \bar{Q}_1\right),$$

are sixth order integrals of motion of $\mathcal{H}'$, and that the set $\mathcal{I}'_1$, $\mathcal{I}'_2$ (1.3) and Q is formed by three functionally independent functions. Alternatively, the six-th order integral Q can be written as

$$Q = \left(p_1^2 + 2\delta q_1^2\right)^2 \left(p_2^2 + 8\delta q_2^2\right) - 16\delta \left(2\delta q_1^2 q_2 + (q_1 p_2 - q_2 p_1) p_1\right)^2 .$$

Now, the second order integral $\mathcal{L}'$ (1.3) arises through the following relation

$$\mathcal{L}'^2 = \frac{8{\mathcal{I}'_1}^2 \mathcal{I}'_2 - Q}{16\delta}.$$

Remarkably enough, for the curved superintegrable Hamiltonian $\mathcal{H}'_\kappa$ (4.3) (in Beltrami variables), the corresponding six-th order integral Q_κ can also be found, namely,

$$Q_\kappa = \left(J_{01}^2 + \kappa J_{12}^2 + 2\delta \, \frac{q_1^2(1+\kappa q_2^2)}{(1-\kappa q_2^2)^2} \right)^2 \left(J_{02}^2 + \frac{8\delta q_2^2}{(1-\kappa q_2^2)^2} \right) - 16\delta \left(\frac{2\delta q_1^2 q_2}{(1-\kappa q_2^2)^2} + J_{12} J_{01} \right)^2 ,$$

where J_{ij} are the generators (2.7). Moreover, the relationship with the second order integral $\mathcal{L}'_\kappa$ (4.3) remains in the same "flat" form

$$\mathcal{L}'^2_\kappa = \frac{8\mathcal{I}'^2_{1,\kappa}\, \mathcal{I}'_{2,\kappa} - Q_\kappa}{16\delta}.$$

Therefore these latter expressions seem to point out that the construction of the $\mathbf{S}^2$ and $\mathbf{H}^2$ analogues of some other Euclidean commensurate oscillators could be feasible.

Secondly, another interesting open problem would be the construction of the constant curvature analogues of some integrable Hénon–Heiles systems (see [24] and references therein) that can be written as particular cases of the multiparametric family

$$\mathcal{H} = \frac{1}{2}(p_1^2 + p_2^2) + \delta q_1^2 + (\delta + \Omega) q_2^2 + \alpha \left(q_1^2 q_2 + \beta q_2^3 \right) + \frac{\lambda_1}{q_1^2},$$

in which the only known integrable cases are the Sawada–Kotera ($\beta = 1/3, \Omega = 0$), the KdV ($\beta = 2, \Omega$ arbitrary) and the Kaup–Kupershmidt ($\beta = 16/3, \Omega = 15\delta$) systems. The results here presented would be a very reasonable starting point, that should be completed with appropriate curved analogues of the cubic Hénon–Heiles term.

Finally, the explicit solution of the Schrödinger equation associated to $\mathcal{H}_\kappa$ and $\mathcal{H}'_\kappa$ should be addressed, as well as the analysis of the quantum nonlinear dynamics generated by these new class of integrable nonlinear quantum models by following, e.g., the approaches presented in [25, 26, 27]. Work on all these lines is in progress.

Acknowledgments

This work was partially supported by the Spanish MINECO under grants MTM2010-18556 and AIC-D-2011-0711 (MINECO-INFN).

References

[1] J. Fris, V. Mandrosov, Y.A. Smorodinsky, M. Uhlir, P. Winternitz, On higher symmetries in quantum mechanics, Phys. Lett. 16 (1965) 354–6.

[2] N.W. Evans, Superintegrability of the Winternitz system, Phys. Lett. A 147 (1990) 483–6.

[3] N.W. Evans, Group theory of the Smorodinsky–Winternitz system, J. Math. Phys. 32 (1991) 3369–75.

[4] C. Grosche, G.S. Pogosyan, A.N. Sissakian, Path integral discussion for Smorodinsky–Winternitz potentials 1. Two- and three-dimensional Euclidean space, Fortschr. Phys. 43 (1995) 453–521.

[5] J.M. Jauch, E.L. Hill, On the problem of degeneracy in quantum mechanics, Phys. Rev. 57 (1940) 641–5.

[6] M.A. Rodríguez, P. Tempesta, P. Winternitz, Reduction of superintegrable systems: The anisotropic harmonic oscillator, Phys. Rev. E 78 (2008) 046608 (6pp).

[7] P.W. Higgs, Dynamical symmetries in a spherical geometry I, J. Phys. A: Math. Gen. 12 (1979) 309–23.

[8] H.I. Leemon, Dynamical symmetries in a spherical geometry II, J. Phys. A: Math. Gen. 12 (1979) 489–501.

[9] C. Grosche, G.S. Pogosyan, A.N. Sissakian, Path integral discussion for Smorodinsky–Winternitz potentials 2. The two- and three-dimensional sphere, Fortschr. Phys. 43 (1995) 523–63.

[10] E.G. Kalnins, W.Jr. Miller, G.S. Pogosyan, Superintegrability of the two-dimensional hyperboloid, J. Math. Phys. 38 (1997) 5416–33.

[11] M.F. Rañada, M. Santander, Superintegrable systems on the two-dimensional sphere S^2 and the hyperbolic plane H^2, J. Math. Phys. 40 (1999) 5026–57.

[12] A. Ballesteros, F.J. Herranz, M. Santander, T. Sanz-Gil, Maximal superintegrability on N-dimensional curved spaces, J. Phys. A: Math. Gen. 36 (2003) L93–9.

[13] A. Ballesteros, F.J. Herranz, Superintegrability on three-dimensional Riemannian and relativistic spaces of constant curvature, SIGMA 2 (2006) 010 (22pp).

[14] A. Ballesteros, A. Enciso, F.J. Herranz, O. Ragnisco, Superintegrability on N-dimensional curved spaces: Central potentials, centrifugal terms and monopoles, Ann. Phys. 324 (2009) 1219–33.

[15] F.J. Herranz, M. Santander, Conformal symmetries of spacetimes, J. Phys. A: Math. Gen. 35 (2002) 6601–18.

[16] F.J. Herranz , R. Ortega, M. Santander, Trigonometry of spacetimes: A new self-dual approach to a curvature/signature (in)dependent trigonometry, J. Phys. A: Math. Gen. 33 (2000) 4525–51.

[17] A. Ballesteros, F.J. Herranz, Maximal superintegrability of the generalized Kepler–Coulomb system on N-dimensional curved spaces, J. Phys. A: Math. Theor. 42 (2009) 245203 (12pp).

[18] E.G. Kalnins, S. Benenti, W.Jr. Miller, Integrability, Stäckel spaces, and rational potentials, J. Math. Phys. 38 (1997) 2345–65.

[19] P. Saksida, Integrable anharmonic oscillators on spheres and hyperbolic spaces, Nonlinearity 14 (2001) 977–94.

[20] M.F. Rañada, M. Santander, On some properties of harmonic oscillator on spaces of constant curvature, Rep. Math. Phys. 49 (2002) 335–43.

[21] M.F. Rañada, M. Santander, On harmonic oscillators on the two-dimensional sphere S^2 and the hyperbolic plane H^2, J. Math. Phys. 43 (2002) 431–51.

[22] M.F. Rañada, M. Santander, On harmonic oscillators on the two-dimensional sphere S^2 and the hyperbolic plane H^2 II, J. Math. Phys. 44 (2003) 2149-67.

[23] F.J. Herranz, A. Ballesteros, M. Santander, T. Sanz-Gil, Maximally superintegrable Smorodinsky–Winternitz systems on the N-dimensional sphere and hyperbolic spaces, in: P. Tempesta et al. (Eds.), Superintegrability in Classical and Quantum Systems, CRM Proceedings and Lecture Notes vol. 37, AMS, Providence RI, 2004, pp. 75–89.

[24] A. Ballesteros, A. Blasco, Integrable Hénon–Heiles Hamiltonians: A Poisson algebra approach, Ann. Phys. 325 (2010) 2787–99.

[25] J.F. Cariñena, M.F. Rañada, M. Santander, A quantum exactly solvable non-linear oscillator with quasi-harmonic behaviour, Ann. Phys. 322 (2007) 434–59.

[26] J.F. Cariñena, M.F. Rañada, M. Santander, The quantum harmonic oscillator on the sphere and the hyperbolic plane, Ann. Phys. 322 (2007) 2249–78.

[27] A. Ballesteros, A. Enciso, F.J. Herranz, O. Ragnisco, D. Riglioni, Quantum mechanics on spaces of nonconstant curvature: The oscillator problem and superintegrability, Ann. Phys. 326 (2011) 2053–73.